arXiv:hep-ph/9804360v1 22 Apr 1998

QUARKONIUM MASS SPLITTING REVISITED: EFFECTS OF CLOSED MESONIC CHANNELS

M.Shmatikov

Russian Research Center "Kurchatov Institute",
123182 Moscow, Russia

Abstract

Modifications of the mass spectrum the quarkonium induced by its virtual dissociation into a pair of heavy mesons is considered. Coupling between quark and mesonic channels results in noticeable corrections to spin-dependent mass splitting. In particular, the observable hierarchy of mass splittings in the χ_c, χ_b and χ'_b multiplets is reproduced.

Quarkonium is the testing ground for various theoretical approaches and, especially, of potential models. In particular, analysis of the quarkonium mass spectrum is used for investigating properties of the confining potential. Phenomenological potentials were reasonably successful in describing gross features of the quarkonia spectra: the level spacing and ordering . More subtle details of the mass spectra, in particular, spin-dependent mass splitting still remains a challenging problem. Time-honored approach to the analysis of spin-dependent effects in quarkonia consists in introducing an interquark *QED-type* potential but with more generic vertex structure [1]:

$$V = \Gamma_i \tilde{V}(q^2) \Gamma_i, \tag{1}$$

where $\tilde{V}(q^2)$ is the propagator of the exchanged particle and Γ_i is the coupling vertex (equal e.g. to 1 or γ^μ in the case of scalar and vector-type coupling respectively) . In the nonrelativistic limit the expansion of (1) in the inverse powers of the heavy-quark mass yields the sum of central and spin-dependent components

$$V = V_c + \left[(\vec{S}_1 + \vec{S}_2)\, \vec{L} \right] V_{LO} + S_{12} \cdot V_T + \left(\vec{S}_1 \cdot \vec{S}_2 \right) V_{SS} \tag{2}$$

with V_{LO}, V_T, V_{SS} standing for the spin-orbit, tensor and spin-spin potentials respectively. In eq.(2) $\vec{S}_{1,2}$ are the quarks' spins, $\vec{L}$ is their orbital momentum and $S_{12} = 12(\vec{S}_1 \cdot \hat{r})(\vec{S}_2 \cdot \hat{r}) - 4(\vec{S}_1 \cdot \vec{S}_2)$. The spin-structure of the interquark potential (2) can be translated in the mass splitting of the triplet P states as follows:

$$\begin{aligned} m(^3P_0) &= \quad \bar{m} - 2a - 4b \\ m(^3P_1) &= \quad \bar{m} - a + 2b \\ m(^3P_2) &= \quad \bar{m} + a - 2b/5 \end{aligned}$$

where $\bar{m}$ is the c.o.g. mass of the triplet (weighted with $2J + 1$ factor), a and b are the averaged values of the V_{LO} and V_T potential components respectively. The V_{SS} component of the potential controls mass splitting between the triplet and singlet P states. The combined action of spin-orbit and tensor forces is parameterized in the form of the ratio

$$R = \frac{m(^3P_2) - m(^3P_1)}{m(^3P_1) - m(^3P_0)} \tag{3}$$

The masses of the ^3P_J charmonium states as determined by Crystal Ball [2] yield

$$R(\chi_c) = 0.478 \pm 0.01 \tag{4}$$

with corresponding ratios for 1^3P_J and 2^3P_J states of the $\bar{b}b$-quarkonium reading respectively [2]

$$R(\chi_b) = 0.664 \pm 0.038 \tag{5}$$

and

$$R(\chi_b') = 0.576 \pm 0.014 \tag{6}$$

Note the hierarchy of R's as given by the measured quarkonia masses:

$$R^{\rm exp}(\chi_b) \gtrsim R^{\rm exp}(\chi_b') > R^{\rm exp}(\chi_c) \tag{7}$$

At the same time potential models involving the sum of the the scalar (confining) and vector (residual one-gluon exchange) potentials (*cf* (1)) yielded controversial results predicting $R(\chi_b) > R(\chi_c)$ and $R(\chi_b')$ slightly larger or, at best, approximately equal to $R(\chi_b)$ in the qualitative disagreement with (7) (see [3] for the details):

$$R(\chi_b') \gtrsim R(\chi_b) > R(\chi_c) \tag{8}$$

Observed disagreement hints at the existence of additional effects. A solution to the problem was sought by introducing some unconventional interactions, in particular, a pseudoscalar component of the quark-antiquark potential with $\Gamma = \gamma^5$ in (1) [4]. Another issue might be an account of relativistic effects which are more pronounced in $\bar{c}c$ as compared to their $\bar{b}b$ counterparts. Currently no convincing solution to the problem in the framework of the potential-based approach seems to exist. The controversy casts doubts on the universally adopted conclusion about scalar Lorentz-transformation properties of the confining potential.

In the present paper we consider an alternative mechanism affecting mass-splitting in the quarkonia systems which might give an insight to the solution of the challenging problem. It is coupling of quarkonia states to pairs of heavy quark-mesons $H\bar{H}$. Investigation of the effects emerging from such coupling was pioneered by [5], where a unified description of the heavy quark-antiquark binding and of the quarkonium coupling to heavy mesons was given. However, the calculations made based heavily on the specific form of the confining potential. The latter was assumed to have Lorentz-vector transformation properties in disagreement with the current experimental data on both the spectroscopy of $\bar{c}c$ states and on their decay amplitudes [6]. The latter circumstance substantiates revisiting effects which emerge from coupling between quarkonia states and mesonic channels.

Coupling of quarkonium ($Q\bar{Q}$) bound states to heavy-meson pairs ($H\bar{H}$) results in the mass shift of the former because of the virtual dissociation

$$Q\bar{Q} \to (Q\bar{q}) + (\bar{Q}q) \to Q\bar{Q} \tag{9}$$

The mass shift emerges as a matter of fact due to an interaction between heavy $Q\bar{Q}$ and (virtual) light $q\bar{q}$ quark pairs. The situation, however, is somewhat tricky. Indeed, in the *adiabatic* approximation the presence of the light-quark pairs does not affect (at least) the long-range part of the potential operating between heavy quark and antiquark. More precisely, its linear form is maintained, and only the tension is altered. The latter, however, is a phenomenological parameter determined from the experimental data implying that in the $m_Q \to \infty$ limit coupling to mesonic channels does not result in an observable effect [7]. This conclusions refers to the spectrum gross features controlled by the string tension. At the same time spin-dependent mass splitting is scaled by a $1/m_Q$ factor, and non-vanishing effect are possible.

Consider in more detail the mechanism which produces effective mass splitting of the P_J triplet. The lightest $Q\bar{q}$ states (and their antiparticles) materialize in the form of pseudoscalar and vector mesons (denoted hereafter H and H^* respectively). Be H and H^* mass-degenerate, coupling between $Q\bar{Q}$ and $q\bar{q}$ would not result in mass splitting of quarkonium states. However, because of the spin $Q-\bar{q}$ interaction the H^* vector meson proves to be heavier than its pseudoscalar H counterpart. It implies that the mass shift of the quarkonium states is scaled by the width of a $H\bar{H}$ continuous spectrum band up to the effective mass $\gtrsim(m_{H^*} - m_H)$. The contribution coming from $H\bar{H}$ pairs with masses exceeding $m_H + m_{H^*}$ is canceled by similar contribution of $\bar{H}H^*$ and $\bar{H}^*H^*$ heavy-meson states.

A quantum-mechanical consideration of the problem allows making some qualitative conclusions concerning the expected effect. First, quarkonium states, depending on their spin-parity quantum numbers, couple to different $H\bar{H}$ states whereas some of them do not have a $H\bar{H}$ match at all. Indeed, the ${}^3P_0$ and ${}^3P_2$ quarkonium states couple to the $H\bar{H}$ pairs with $J^\pi = 0^+$ and 2^+ respectively, while the ${}^3P_1$ level is uncoupled from the (pseudoscalar) mesonic sector. The ${}^1P_1$ state does not couple to $H\bar{H}$ mesons either. Selective coupling results in different mass shifts of the $Q\bar{Q}$ states, and, eventually, in their mass splitting. Second, the $H\bar{H}$ continuous spectrum lies above the quarkonium bound states. Spin-independent potentials reproduce the gross features of the spectrum accurately enough to assume that the spin-dependent forces may be treated as a perturbation. The second-order perturbation in energy [8] equals

$$E_n^{(2)} = \sum_{m \neq n} \frac{|V_{nm}|^2}{E_n^{(0)} - E_m^{(0)}} \tag{10}$$

where V_{nm} is a perturbing interaction and $E_n^{(0)}$ is the (unperturbed) energy of the n-th state. Eq.(10) shows that the mass of the lowest state E_0 (i.e. the quarkonium) decreases because of channel coupling (since $E_0^{(0)} < E_m^{(0)}$).

Combining these observations we can conclude that the masses of both the ${}^3P_0$ and ${}^3P_2$ states will diminish because of their coupling to mesonic $H\bar{H}$ states, whereas the mass of the ${}^3P_1$ state will remain unaffected. Inspection of (3) shows immediately that the considered mechanism results in the decrease of the R ratio.

Next observation is that the magnitude of the effect depends on the width of the energy gap separating quarkonium states from the continuous $H\bar{H}$ spectrum: the smaller is the gap, the more pronounced will be the mass shift. We infer then that the $R(\chi_b)$ ratio (1^3P_J bottomonium states) will be affected less than the $R(\chi_b')$ ratio (2^3P_J bottomonium states). Comparing these *qualitative* predictions to the experimental data (7) we conclude that the considered mechanism (coupling between quarkonium and mesonic states) gives a welcome trend to the modified theoretical results.

We proceed now to quantitative estimates of the effect. To this end a dynamical model describing the coupling mechanism under consideration is to developed. We treat quarkonium ($Q\bar{Q}$) and mesonic ($H\bar{H}$) states as the components of a two-channel system labeling them channels 1 and 2 respectively. Then the Schrödinger equations describing the dynamics of the system for each J^π state can be written as follows

$$\begin{array}{llll} T_1\phi_1 + & V_{11}\phi_1 + & V_{12}\phi_2 = & E_1\phi_1 \\ T_2\phi_2 + & & V_{21}\phi_1 = & E_2\phi_2 \end{array} \qquad (11)$$

Here T is the kinetic energy in the corresponding channels and V is the interaction potential. The V_{11} component is a (linear + Coulombic) potential operating between heavy quarks. Interaction between H-mesons is neglected: near-threshold phenomena under consideration involve small relative momenta and, hence, large inter-particle distances. At the same time, the long-range interaction - $\pi-$meson exchange - between two pseudoscalar particles is absent. Anyway, this approximation does not alter the qualitative conclusions made above. Finally, E_1 and E_2 are the channel energies, differing by the mass gap in the corresponding state: $E_2 - E_1 = 2m_H - m_{Q\bar{Q}}$. The non-diagonal component of the potential V_{12} ($= V_{21}$) describes the dynamics of the quarkonium dissociation $(Q\bar{Q}) \rightarrow H\bar{H}$, or, stated differently, the dynamics of the $q\bar{q}$ production in the field of heavy quarks. Forbidding complexity of the problem necessitates introducing some simplifying approximations.

Given the large mass difference between heavy and light quarks, we make use of the adiabatic approximation. The latter molds in the following assumptions. First, the position of the heavy quarks is not affected by the process of the $q\bar{q}$ production. Second, the combined spin of heavy quarks in a pair of heavy mesons is equal to zero. It implies that in all of the transitions considered the spin of the $Q\bar{Q}$ pair is subject to one and the same change from $S = 1$ to $S = 0$. Finally, c.o.g. of a heavy meson coincides with good accuracy with the position of the heavy meson residing in it. All the assumptions made above can be expressed concisely in the following form of the non-diagonal component:

$$V_{12} = \beta \, \delta \, (\vec{r} - \vec{r}\,') \tag{12}$$

where β is a coupling strength constant, $\vec{r}$ and $\vec{r}\,'$ are the separation between heavy quarks prior and after the $q\bar{q}$ production (i.e. within the quarkonium state and in the $H\bar{H}$ pair respectively). The light-quark spin operator and coordinate variables are assumed to be present and are dropped for notational brevity.

In the framework of the model the strength constant β can be determined without going into details of the complicated mechanism of the light-quark pair production. The same coupling between channels governs decays of the quarkonium states lying *above* the $H\bar{H}$ threshold, i.e. $\psi(3770)$ and $\Upsilon(10580)$ in the case of charmonium and bottomonium respectively. They are known to decay predominantly into the $D\bar{D}$ and $B\bar{B}$ pairs with the widths equal to 83.9±2.4 and 21±4 MeV respectively. The quantum numbers of the bottomonium state are known ($J^{\pi} = 1^{-}$) translating into the 4^3S_1 state of the $b\bar{b}$ pair. The $\psi(3770)$ quantum numbers are not determined; following the potential model prescriptions we adopt that it is the 1^3D_1 state of the $c\bar{c}$ pair [5, 9]. Using the channel-coupling potential (12) we calculate the value of the β strength coupling constant. It proves to be equal to 153 MeV and 114 MeV for the charmonium and bottomonium systems respectively. The agreement between two values of β is reasonably good signaling the validity of the approximations made. Indeed, the paradigm of the non-perturbative QCD assumes the light-quark pair production as a result of the string break-up. Flavor-blindness of the confinement forces (materialized as a string) prompts then that in the infinite heavy-quark mass limit the values of β for two types of the quarkonium should the same. Given possible $1/m_Q$ corrections, the β values may be considered as coinciding in line with the theoretical expectation.

Having determined the potential V_{12} connecting the quarkonium and the mesonic channels we proceed to calculating the matrix element of the transition between them. The matrix element controls the value of the mass shift (see (10)). Within the approximations made it reduces to the overlap integral of the quarkonium wave function $\psi^{Q\bar{Q}}$ and of the wave function $\psi_{\nu}^{H\bar{H}}$ of two non-interacting heavy mesons:

$$V_{\nu} = \beta \int \psi^{Q\bar{Q}}(\vec{r}) \cdot \psi_{\nu}^{H\bar{H}}(\vec{r}) \, d\vec{r} \tag{13}$$

The subscript ν labels the states of the $H\bar{H}$ continuous spectrum, to this end we use the c.m.s. momentum of heavy mesons ($\nu \equiv \vec{k}$). Then the expression for the quarkonium-state mass shift (10) can be cast in the form

$$\Delta m_{Q\bar{Q}} = \int \frac{\left|V_{\vec{k}}\right|^2}{m_{Q\bar{Q}} - m_{H\bar{H}}}\, d\vec{k} \tag{14}$$

where $m_{H\vec{H}} = 2m_H + k^2/m_H$. Formal convergence of the integral is ensured by (slow) decrease of the $V_{\vec{k}}$ matrix element due to diminishing overlap between the quarkonium bound-state wave function and the oscillating wave function of two heavy mesons in the continuous spectrum. However, the integration over $\vec{k}$ which may be translated in the integration over the mass of the $H\bar{H}$ pair is to be cut-off at much smaller $m_{\bar{H}H}$ values. Indeed, the considered mechanism produces mass splitting of the ^3P_J states because of the mass difference between the pseudoscalar H-meson and its vector counterpart. We incur then that the scale of integration range is set by the $m_{H^*} - m_H$ mass difference.

Results of numerical calculations confirm the qualitative conclusions drawn above. To investigate sensitivity of the effect we cut off the integration in (14) at some μ ($\mu \geq 2m_H$). Dependence of the R-ratio for the $2P$ and $1P$ states of the bottomonium and for the $1P$ states of the charmonium are displayed in fig.1 as a function of the reduced mass band $r = (\mu/2 - m_H)/(m_{H^*} - m_H)$. As the initial values for R's we used those calculated in a potential model [10] (using other predictions of potential models does not change qualitatively the obtained results). Note the fast onset of the $R(2P) \lesssim R(1P)$ regime: already for $r \gtrsim 0.2$ the hierarchy of the R values corresponds to the experimental observations (7) and it maintains with further increase of r. The considered mechanism does not yield quantitative agreement with the experimental data: the overall set of data is shifted downwards with respect to experimental values (shown by error bars). A possible explanation may be that parameters of the spin-dependent potential determining the "initial" values of R were varied already in an attempt to reproduce experimentally observed regularities. It looks plausible that the consistent description of spin-dependent mass splitting in the quarkonium systems requires the account of both the spin-dependent part of the $Q - \bar{Q}$ potential and of the bound-state coupling to mesonic channels. The investigation of the problem following these lines is in progress.

Summarizing, the process of (virtual) light-quark pair production considerably affects masses of the quarkonium states. Account of the emerging mass splitting of the ${}^3P_J$ triplet allows to reproduce qualitative features of the experimentally observed quarkonium level spacing.

The author is indebted to F.Lev and S.Romanov for helpful comments.

References

[1] J.F.Donogue, E.Golowich and B.R.Holstein. Dynamics of the Standard Model, *Cambridge University Press*, 1992;

[2] Particle Data Group, Phys.Rev. **D54** (1996) 1;

[3] K.K.Seth, In: Proc. of the Third Workshop on the Tau-Charm Factory. Mirabelle, Spain, June 1993. Eds. J.Kirkby and R.Kirkby. *Editions Frontieres*, p. 461;

[4] J.Lee-Franzini and P.J.Franzini, *ibid.* p. 447;

[5] E.Eichten *et al.* Phys.Rev. **D17** (1978) 3090; *ibid.* **D21** (1980) 203;

[6] T.Barnes, In: Proc. of Hadron '95, The Sixth Intern. Conf. on Hadron Spectroscopy, Manchester, UK, July 1995. Eds. M.C.Birse, G.D.Lafferty and J.A.McGovern. *World Scientific*, p. 33;

[7] P.Geiger and N.Isgur, Phys.Rev. **D41** (1990) 1595;

[8] L.D.Landau and E.M.Lifshitz, Quantum Mechanics: Nonrelativistic Theory, 3rd ed., *Pergamon Press*, Oxford - New York, 1977;

[9] S.Godfrey and N.Isgur, Phys.Rev. **D32** (1985) 189;

[10] S.N.Gupta, S.F.Radford and W.W.Repko, Phys.Rev. **D26** (1982) 3305; *ibid.* **D34** (1986) 201


R_χ
r
0.8
0.7
0.6
0.5
0.4
0.3
0.2
0
0.2
0.4
0.6
0.8
1
$R_{\chi_b}^{\rm exp}$
$R_{\chi_b'}^{\rm exp}$
$R_{\chi_c}^{\rm exp}$
R_{χ_b}
$R_{\chi_b'}$
R_{χ_c}